

What is Claimed is:

1. A method of preventing and treating a skin disorder comprising the steps of:  
providing a composition comprising a quantity of *Agaricus blazei* Murill; and  
administering an effective amount of said composition to a subject.
2. A method as described in claim 1, wherein the composition is administered orally.
3. A method as described in claim 1, wherein the composition is formulated in a liquid solution.
4. A method as described in claim 1, wherein the composition is in a form selected from the group consisting of a powder, crystalline and granule.
5. A method as described in claim 1, wherein the composition is administered topically.
6. A method as described in claim 1, wherein the composition is formulated as one of the group consisting of: a cream, ointment, gel, cosmetic lotion and a shampoo.
7. A method as described in claim 1, wherein the *Agaricus blazei* Murill is present in a concentration of at least about 10mg per unit composition.
8. A method as described in claim 1, wherein the *Agaricus blazei* Murill is present in a concentration of between about 10mg and 600mg per unit composition.
9. A method as described in claim 1, wherein the *Agaricus blazei* Murill is present in a concentration of about 100mg per unit composition.
10. A method as described in claim 1, wherein the *Agaricus blazei* Murill is an extract obtained by water extraction.
11. A method as described in claim 1, wherein the *Agaricus blazei* Murill is an extract obtained according to the following steps:

1

providing whole or particulate *Agaricus blazei* Murill;  
 extracting the whole or particulate *Agaricus blazei* Murill in hot water to obtain a residue;  
 extracting the residue in an aqueous solution of ammonium oxalate to obtain an extract;  
 5 decomposing the extract in hydrochloric acid;  
 gel permeating the extract; and  
 purifying the extract with affinity chromatography.

5

10

12. A method as described in claim 10 or 11, wherein the extract is at least one substance selected from the group consisting of: a polysaccharide-glucan, a steroid, a dietary fiber; linoleic acid; ergosterol; nicotinic acid amide; benzoic acid and beta glucans.

15

13. A method as described in claim 1, wherein the composition is formulated as a pill or capsule.

20

14. A method as described in claim 1, wherein the composition further comprises other active or inactive ingredients.

25

15. A method as described in claim 1, wherein the skin disorder is one selected from the group consisting of: neoplasms, melanoma, multicentric primary skin malignancy, and viral skin disease.

30

16. A method as described in claim 1, wherein the skin disorder is giant congenital melanocytic nevi.

35

17. A method as described in claim 1, wherein the skin disorder is either a basal cell or squamous cell cancer.

18. A method as described in claim 1, wherein the skin disorder is either mycosis fungoides or Kaposi's sarcoma.

19. A method as described in claim 1, wherein the skin disorder is either viral warts or molluscum contagiosum.

1

20. A method as described in claim 1, wherein the *Agaricus blazei* Murill is an extract and is extracted at a ratio of about one part *Agaricus blazei* to about 2 to 5 parts aqueous solution.

5

21. A method as described in claim 1, wherein the composition contains at least about 0.01% *Agaricus blazei* Murill by volume.

10

22. A method as described in claim 1, wherein the composition contains at least about 0.05% *Agaricus blazei* Murill by volume.

23. A method as described in claim 1, wherein the composition contains at least about 0.05% to about 20% *Agaricus blazei* Murill by volume.

15

24. A method as described in claim 1, wherein the subject is a domesticated or livestock animal.

25. A method as described in claim 1, wherein the subject is a human being.

20

26. A method as described in claim 1, wherein the composition is administered at least once daily for between about 10 to about 30 days.

27. A method as described in claim 1, wherein an interval of between about 1 and about 10 days is provided between administrations of the compound.

25

28. A method as described in claim 1, wherein the composition is formulated into either an animal food or a dietary supplement for animals

29. A method as described in claim 1, wherein the *Agaricus blazei* Murill is in a form selected from the group consisting of: whole, particulate and extract thereof.

30

30. A method as described in claim 1, wherein the composition comprises *Agaricus blazei* Murill from all four stage of the mushroom lifecycle.

35

31. A method of preventing and treating an autoimmune disorder comprising the steps of:

1

providing a composition comprising *Agaricus blazei* Murill; and  
administering an effective amount of said composition to a subject.

5

32. A method as described in claim 31, wherein the composition is administered orally.

10

33. A method as described in claim 31, wherein the composition is formulated in a liquid solution.

34. A method as described in claim 31, wherein the composition is in a form selected from the group consisting of a powder, crystalline and granule.

15

35. A method as described in claim 31, wherein the composition is administered topically.

36. A method as described in claim 31, wherein the composition is formulated as one of the group consisting of: a cream, ointment, gel, cosmetic lotion and a shampoo.

20

37. A method as described in claim 31, wherein the *Agaricus blazei* Murill is present in a concentration of at least about 10mg per unit composition.

25

38. A method as described in claim 31, wherein the *Agaricus blazei* Murill is present in a concentration of between about 10mg and 600mg per unit composition.

39. A method as described in claim 31, wherein the *Agaricus blazei* Murill is present in a concentration of about 100mg per unit composition.

30

40. A method as described in claim 31, wherein the *Agaricus blazei* Murill is an extract obtained by water extraction.

41. A method as described in claim 31, wherein the *Agaricus blazei* Murill is an extract obtained according to the following steps:

35

providing whole or particulate *Agaricus blazei* Murill;  
extracting the whole or particulate *Agaricus blazei* Murill in hot water to obtain a residue;

extracting the residue in an aqueous solution of ammonium oxalate to obtain an extract;  
decomposing the extract in hydrochloric acid;  
gel permeating the extract; and  
purifying the extract with affinity chromatography.

42. A method as described in claims 40 or 41, wherein the extract is at least one substance selected from the group consisting of: a polysaccharide-glucan, a steroid, a dietary fiber; linoleic acid; ergosterol; nicotinic acid amide; benzoic acid and beta glucans.

43. A method as described in claim 31, wherein the composition is formulated as a pill or capsule.

44. A method as described in claim 31, wherein the composition further comprises other active or inactive ingredients.

45. A method as described in claim 31, wherein the autoimmune disorder is one selected from the group consisting of: intravascular disorders, immune disorders and neuromuscular disorders.

46. A method as described in claim 31, wherein the autoimmune disorder is selected from the group consisting of: hemolytic anemia, thyroid goiter, Hashimoto's disease, pernicious anemia, type I and II diabetes mellitus and chronic hepatitis.

47. A method as described in claim 31, wherein the autoimmune disorder is selected from the group consisting of: glomerulonephritis, systemic lupus erythematosus and Sjogren's disease.

48. A method as described in claim 31, wherein the autoimmune disorder is selected from the group consisting of: myasthenia gravis and Graves' disease.

49. A method as described in claim 31, wherein the autoimmune disorder is a disease selected from the group consisting of: rheumatoid arthritis, multiple sclerosis and ulcerative colitis.

1

50. A method as described in claim 31, wherein the *Agaricus blazei* Murill is an extract and is extracted at a ratio of about one part *Agaricus blazei* to about 2 to 5 parts aqueous solution.

5

51. A method as described in claim 31, wherein the *Agaricus blazei* Murill is an extract and is extracted at a ratio of about one part *Agaricus blazei* to between about 25 to 100 parts aqueous solution.

10

52. A method as described in claim 31, wherein the composition contains at least about 0.01% *Agaricus blazei* Murill by volume.

15

53. A method as described in claim 31, wherein the composition contains at least about 0.05% *Agaricus blazei* Murill by volume.

54. A method as described in claim 31, wherein the composition contains at least about 0.05% to about 20% *Agaricus blazei* Murill by volume.

20

55. A method as described in claim 31, wherein the subject is an animal.

56. A method as described in claim 31, wherein the subject is a human being.

25

57. A method as described in claim 31, wherein the composition is administered at least once daily for between about 10 to about 30 days.

58. A method as described in claim 31, wherein an interval of between about 1 and about 10 days is provided between administrations of the compound.

30

59. A method as described in claim 31, wherein the composition is formulated into either an animal food or a dietary supplement for animals

35

60. A method as described in claim 31, wherein the *Agaricus blazei* Murill is in a form selected from the group consisting of: whole, particulate and extract thereof.

1

61. A method as described in claim 31, wherein the composition comprises *Agaricus blazei* Murill from all four stage of the mushroom lifecycle.

5

62. A method of preventing and treating a skin disorder comprising the steps of:  
providing a composition comprising an *Agaricus blazei* Murill; and  
administering at least about 10mg of said composition per day to a subject over a period  
of between about 10 and 30 days.

10

63. A method as described in claim 62, wherein an interval of between about 1 and  
about 10 days is provided between administrations of the compound.

15

64. A method as described in claim 62, wherein the composition is administered to  
the subject either orally or topically.

20

65. A method of preventing and treating an autoimmune disorder comprising the steps  
of:  
providing a composition comprising an *Agaricus blazei* Murill; and  
administering at least about 10mg of said composition per day to a subject over a period  
of between about 10 and 30 days.

25

66. A method as described in claim 65, wherein an interval of between about 1 and  
about 10 days is provided between administrations of the compound.

30

67. A method as described in claim 65, wherein the composition is administered to  
the subject either orally or topically.

35